ELSEVIER

Environmental Pollution 139 (2006) I–IV

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 139

Number 1 2006

1 **Interactions of mycorrhizal fungi with *Pteris vittata* (As hyperaccumulator) in As-contaminated soils**
H.M. Leung, Z.H. Ye, M.H. Wong

Indigenous mycorrhizal fungi play an important role in As tolerance.

9 **Polycyclic aromatic hydrocarbons in suspended particulate matter and sediments from the Pearl River Estuary and adjacent coastal areas, China**
X.-J. Luo, S.-J. Chen, B.-X. Mai, Q.-S. Yang, G.-Y. Sheng, J.-M. Fu

PAH were determined in suspended particulate matter and sediments from Pearl River Estuary.

21 **Modelling the extra and intracellular uptake and discharge of heavy metals in *Fontinalis antipyretica* transplanted along a heavy metal and pH contamination gradient**
J.A. Fernández, M.D. Vázquez, J. López, A. Carballeira

The kinetics of uptake and discharge of heavy metals, in different cellular locations, were studied in transplanted aquatic mosses.

32 **Bioconcentration of atrazine and chlorophenols into roots and shoots of rice seedlings**
Y.-H. Su, Y.-G. Zhu

Uptakes of o-chlorophenol, 2,4-dichlorophenol, and atrazine at various levels from nutrient solution by roots and shoots of rice seedlings were investigated using a partition-limited model.

40 **Soil amendments reduce trace element solubility in a contaminated soil and allow regrowth of natural vegetation**
E. Madejón, A.P. de Mora, E. Felipe, P. Burgos, F. Cabrera

Soil amendments affect soil chemistry and allow revegetation of soils contaminated by trace elements.

53 **Evidence for potential impacts of ozone on *Pinus cembra* L. at mountain sites in Europe: An overview**
G. Wieser, W.J. Manning, M. Tausz, A. Bytnerowicz

An overview of ozone effects on Pinus cembra an important European coniferous tree species in the timberline ecotone.

59 **Perchlorate in fish from a contaminated site in east-central Texas**
C. Theodorakis, J. Rinchard, T. Anderson, F. Liu, J.-W. Park, F. Costa, L. McDaniel, R. Kendall, A. Waters

In perchlorate-contaminated lakes and streams, perchlorate is detected infrequently in fish heads, fillets, and whole bodies, but may be detected more often depending on species and seasonal trends, and always at concentrations higher in the fish than in the water.

70 **A comparative assessment of heavy metal accumulation in soft parts and byssus of mussels from subarctic, temperate, subtropical and tropical marine environments**
P. Szefer, S.W. Fowler, K. Ikuta, F.P. Osuna, A.A. Ali, B.-S. Kim, H.M. Fernandes, M.-J. Belzunce, B. Guterstam, H. Kunzendorf, M. Wołowicz, H. Hummel, M. Deslous-Paoli

Byssus of mytilids, like soft tissues can be used as efficient biomonitor for heavy metals in the marine environment.

79 **Tetrodotoxin prevents copper-induced bradycardia in gastropod limpets**
G. Bini, A.M. Pugliese, G. Pepeu, G. Chelazzi

In limpets, copper-induced bradycardia involves the activation of an extrinsic neuronal pathway.

86 **Release of trace metals, sulfate and complexed cyanide from soils contaminated with gas-purifier wastes: A microcosm study**
T. Rennert, T. Mansfeldt

The potential release of trace metals, sulfate and cyanide from soils contaminated with gas-purifier wastes is a function of pH and the redox.

95 **Contamination by arsenic and other trace elements in tube-well water and its risk assessment to humans in Hanoi, Vietnam**
T. Agusa, T. Kunito, J. Fujihara, R. Kubota, T.B. Minh, P.T. Kim Trang, H. Iwata, A. Subramanian, P.H. Viet, S. Tanabe

High concentrations of arsenic, manganese and barium were found in tube-well water and human hair in suburban areas of Hanoi, Vietnam.

107 Specific accumulation of organochlorines in human breast milk from Indonesia: Levels, distribution, accumulation kinetics and infant health risk
A. Sudaryanto, T. Kunisue, N. Kajiwara, H. Iwata, T.A. Adibroto, P. Hartono, S. Tanabe

Specific residents were exposed to high levels of DDTs in Indonesia.

118 Effects of teflubenzuron on sediment processing by members of the *Capitella* species-complex
N. Méndez

Capitella sp I reduces its feeding activity at concentrations from 8.4 to 41.8 μg/g dry wt. sediment of the insecticide teflubenzuron.

125 Scale and causes of lead contamination in Chinese tea
W.-Y. Han, F.-J. Zhao, Y.-Z. Shi, L.-F. Ma, J.-Y. Ruan

32% of Chinese tea samples exceeded the national maximum permissible concentration for Pb.

133 Development of an SPME–GC–MS/MS method for the determination of pesticides in rainwater: Laboratory and field experiments
N. Sauret-Szczepanski, P. Mirabel, H. Wortham

Solid-phase microextraction efficiency of pesticides in rainwater was optimized.

143 Water quality dynamics and hydrology in nitrate loaded riparian zones in the Netherlands
M. Hefting, B. Beltman, D. Karssenberg, K. Rebel, M. van Riessen, M. Spijker

Riparian zones reduced nitrate from agricultural lands.

157 Treatment of log yard run-off by irrigation of grass and willows
M. Jonsson, I. Dimitriou, P. Aronsson, T. Elowson

By irrigating willow and couchgrass soil-plant systems with log yard run-off water, TOC, phenols, and phosphorus were reduced with 35% to 96% in the water.

167 Effect of soil characteristics on Cd uptake by the hyperaccumulator *Thlaspi caerulescens*
J. Yanai, F.-J. Zhao, S.P. McGrath, T. Kosaki

Higher total/extractable Cd in soil, lower pH and coarser texture were associated with higher Cd concentration by Thlaspi caerulescens.

176 Use of dolomite phosphate rock (DPR) fertilizers to reduce phosphorus leaching from sandy soil
G.C. Chen, Z.L. He, P.J. Stoffella, X.E. Yang, S. Yu, D. Calvert

Fertilizers developed from dolomite phosphate rock (DPR) reduce phosphorus leaching from sandy soil.

183 Modeling the dynamic changes in concentrations of γ-hexachlorocyclohexane (γ-HCH) in Tianjin region from 1953 to 2020
S. Tao, Y. Yang, H.Y. Cao, W.X. Liu, R.M. Coveney, F.L. Xu, J. Cao, B.G. Li, X.J. Wang, J.Y. Hu, J.Y. Fang

Levels of lindane in various media of Tianjin area decreased significantly during the last decade and will drop more than one order of magnitude in the next 20 years.

Number 2 2006

195 Primary organic pollutants in New Zealand urban aerosol in winter during high PM_{10} episodes
Z. Krivácsy, M. Blazsó, D. Shooter

PM_{10} aerosol concentrations can exceed air quality guidelines during winter in Christchurch, New Zealand.

206 The influence of salinity on metal uptake and effects in the midge *Chironomus maddeni*
J.R. Bidwell, J.R. Gorrie

Elevated porewater salinity significantly influenced uptake of copper, lead and zinc by midge larvae.

214 Ecotoxicological effects of metal pollution in two mollusc species from the Spanish South Atlantic littoral
V. Funes, J. Alhama, J.I. Navas, J. López-Barea, J. Peinado

Oysters adapt to metal pollution while mussels are more sensitive.

224 Seasonal patterns of ascorbate in the needles of Scots Pine (*Pinus sylvestris* L.) trees: Correlation analyses with atmospheric O_3 and NO_2 gas mixing ratios and meteorological parameters
K. Haberer, L. Jaeger, H. Rennenberg

Contents of apoplastic but not symplastic ascorbate correlate significantly with atmospheric ozone concentrations.

232 Radionuclides, heavy metals and fluorine incidence at Tapira phosphate rocks, Brazil, and their industrial (by) products
F.T. da Conceição, D.M. Bonotto

Radionuclides and metals in phosphate fertilizers are not hazardous.

244 Genetic diversity of culturable bacteria in oil-contaminated rhizosphere of *Galega orientalis*
M.M. Jussila, G. Jurgens, K. Lindström, L. Suominen

Bacterial diversity during rhizoremediation in oil-contaminated soil is characterized by a combination of molecular methods.

258 Inorganic $PM_{2.5}$ at a U.S. agricultural site
J.T. Walker, W.P. Robarge, A. Shendrikar, H. Kimball

At this agricultural site, sufficient atmospheric ammonia exists such that inorganic $PM_{2.5}$ formation is limited by the availability of atmospheric acids.

272 Potential application of synchronous fluorescence spectroscopy to determine benzo[*a*]pyrene in soil extracts
G. Hua, K. Killham, I. Singleton

Synchronous fluorescence spectroscopy has potential as a method for confirmation of benzo[a]pyrene in soil extracts.

279 Spatial variation in the degradation rate of the pesticides isoproturon, azoxystrobin and diflufenican in soil and its relationship with chemical and microbial properties
G.D. Bending, S.D. Lincoln, R.N. Edmondson

Spatial variation determines risk assessment for pesticides in soil.

288 Phosphorus fractions and the effect of pH on the phosphorus release of the sediments from different trophic areas in Taihu Lake, China
X. Jin, S. Wang, Y. Pang, F. Chang Wu

Factors affecting internal loading of phosphorus were investigated in a hypereutrophic lake in China.

296 The landsnail *Cepaea nemoralis* regulates internal Cd levels when fed on Cd-enriched stinging nettle (*Urtica dioica*) leaves at low, field-relevant concentrations
M.J.M. Notten, A.J.P. Oosthoek, J. Rozema, R. Aerts

Cd excretion via the mucus plays a substantial role in the regulation of C. nemoralis soft tissue Cd concentrations at low, but field-relevant Cd concentrations in the food.

306 Development of a new biotic index to assess freshwater pollution
J.-G. Jiang

A new biotic index of water quality based on protozoa is described.

318 An in situ postexposure feeding assay with *Carcinus maenas* for estuarine sediment-overlying water toxicity evaluations
S.M. Moreira, M. Moreira-Santos, L. Guilhermino, R. Ribeiro

A new sub-lethal toxicity assay is presented for marine invertebrates.

330 The combined use of chemical and biochemical markers to assess water quality along the Ebro River
R. Lavado, R. Ureña, R. Martin-Skilton, A. Torreblanca, J. del Ramo, D. Raldúa, C. Porte

Fish accumulation of persistent organic pollutants is compared with various biochemical markers.

340 Brominated flame retardants and organochlorine pollutants in aquatic and terrestrial predatory birds of Belgium: Levels, patterns, tissue distribution and condition factors
V.L.B. Jaspers, A. Covaci, S. Voorspoels, T. Dauwe, M. Eens, P. Schepens

Aquatic and terrestrial predatory birds show different accumulation profiles of PBDEs.

353 The assessment of the energetic compound 2,4,6,8,10,12-hexanitro-2,4,6,8,10,12-Hexaazaisowurtzitane (CL-20) degradability in soil
N. Strigul, W. Braida, C. Christodoulatos, W. Balas, S. Nicolich

Biodegradation and plant uptake suggest possible entry of CL-20 into food chain.

362 Colonisation of a Zn, Cd and Pb hyperaccumulator *Thlaspi praecox* Wulfen with indigenous arbuscular mycorrhizal fungal mixture induces changes in heavy metal and nutrient uptake
K. Vogel-Mikuš, P. Pongrac, P. Kump, M. Nečemer, M. Regvar

Colonisation of a Zn, Cd and Pb hyperaccumulator Thlaspi praecox with arbuscular mycorrhizal fungi resulted in improved nutrient and reduced Cd and Zn uptake.

372 Relationships and bioaccumulation of chemical elements in the Baikal seal (*Phoca sibirica*)
T. Ciesielski, M.V. Pastukhov, P. Fodor, Z. Bertenyi, J. Namieśnik, P. Szefer

Biomagnification may be occurring in the fish–seal trophic link.

Number 3 2006

385 Comparative acute toxicity of organic pollutants and reference values for crustaceans. I. Branchiopoda, Copepoda and Ostracoda
F. Sánchez-Bayo

A reference guide for the effects of chemicals on crustaceans.

421 Growth and photosynthetic responses of two pine species (*Pinus koraiensis* and *Pinus rigida*) in a polluted industrial region in Korea
D.S. Choi, M. Kayama, H.O. Jin, C.H. Lee, T. Izuta, T. Koike

Pinus koraiensis seems to be more pollution tolerant than Pinus rigida.

433 Effect of bone char application on Pb bioavailability in a Pb-contaminated soil
S.-B. Chen, Y.-G. Zhu, Y.-B. Ma, G. McKay

Bone char amendments show potential for remediation of Pb-contaminated soils.

440 Afforestation, seasalt episodes and acidification – A paired catchment study in western Norway
T. Larssen, J. Holme

Seasalt episodes cause higher mobilization of toxic aluminum in sites afforested with spruce.

451 Avoidance of low doses of naphthalene by Collembola
L. Boitaud, S. Salmon, C. Bourlette, J.-F. Ponge

PAH avoidance by soil springtails is species-specific and differs among populations of the same species.

455 The influence of single and multiple applications of pyrene on the evolution of pyrene catabolism in soil
C.J.A. Macleod, K.T. Semple

Evolution of catabolic activity in soil is influenced by the concentration and number of applications of organic contaminants.

461 Records of atmospheric delivery of pyrolysis-derived pollutants in recent mountain lake sediments of the Julian Alps (NW Slovenia)
G. Muri, S.G. Wakeham, N.L. Rose

Fluxes of black carbon, polycyclic aromatic hydrocarbons and spheroidal carbonaceous particles were determined in recent remote alpine lake sediments.

469 Genetic toxicity in surface water from Guaíba Hydrographic Region under the influence of industrial, urban and agricultural sewage in the Drosophila Wing-Spot Test
V.S. do Amaral, M. Sinigaglia, M.L. Reguly, H.H.R. de Andrade

Drosophila Wing-Spot Test can be used for detection of environmental mutagenesis.

477 An interdisciplinary physical–chemical approach for characterization of arsenic in a calciner residue dump in Cornwall (UK)
J.T. van Elteren, Z. Šlejkovec, I. Arčon, H.-J. Glass

Arsenic in calciner residue dumps is present as arsenate in crystalline ($Fe_2(As(AsO_4)_3)$) and amorphous or poorly-crystalline (bound to hematite and quartz) structures.

489 Determination of potential sources of PCBs and PBDEs in sediments of the Niagara River
F. Samara, C.W. Tsai, D.S. Aga

Wastewater treatment plant discharges are a main source of PCBs and PBDEs to Niagara River sediments.

498 Stabilization of mercury-containing wastes using sulfide
H. Piao, P.L. Bishop

Sulfide is an effective reagent for the stabilization of mercury when optimum stabilization pH and sulfide dosage are used.

507 Heavy metal speciation and phytotoxic effects of three representative sewage sludges for agricultural uses
I. Walter, F. Martínez, V. Cala

Different sewage sludge processing methods affected the distribution of metals and the seed germination test in different ways.

515 Tradescantia micronucleus test indicates genotoxic potential of traffic emissions in European cities
A. Klumpp, W. Ansel, G. Klumpp, V. Calatayud, J.P. Garrec, S. He, J. Peñuelas, À. Ribas, H. Ro-Poulsen, S. Rasmussen, M.J. Sanz, P. Vergne

The Tradescantia micronucleus test can be used to assess genotoxic potential at urban sites.

523 GC and GC–MS characterization of crude oil transformation in sediments and microbial mat samples after the 1991 oil spill in the Saudi Arabian Gulf coast
T. Garcia de Oteyza, J.O. Grimalt

Cyanobacterial mats inhibit degradation of oil by reducing exposure to the atmosphere and seawater.

532 Lac Dufault sediment core trace metal distribution, bioavailability and toxicity to *Hyalella azteca*
M. Nowierski, D.G. Dixon, U. Borgmann

Toxicity and bioaccumulation tests with sediment cores provide more information on biological effects of metals than surface sediment tests.

541 Effect of copper on the adsorption of *p*-nitrophenol onto soils
Z.-g. Pei, X.-q. Shan, B. Wen, S. Zhang, L. Yan, S.U. Khan

The adsorption of p-nitrophenol onto soils decreased in the presence of copper due to its competition for the adsorption sites.

550 A microcosm test of adaptation and species specific responses to polluted sediments applicable to indigenous chironomids (Diptera)
S. Bahrndorff, J. Ward, V. Pettigrove, A.A. Hoffmann

A field test for adaptation applicable to indigenous chironomids suggests adaptation to metal contaminants in one species but not in other species.

561 Tissues and hair residues and histopathology in wild rats (*Rattus rattus* L.) and Algerian mice (*Mus spretus* Lataste) from an abandoned mine area (Southeast Portugal)
R. Pereira, M.L. Pereira, R. Ribeiro, F. Gonçalves

The bioaccumulation of As and Cd and signs of renal histopathological injury proved the value of Algerian mice as a bioindicator species in the risk assessment of contaminated sites.